



CERTSWARRIOR

Nursing Nurse-Practitioner Nurse Practitioner Exam (NP)

Questions&AnswersPDF

ForMoreInformation:

<https://www.certswarrior.com/>

Features:

- 90DaysFreeUpdates
- 30DaysMoneyBackGuarantee
- InstantDownloadOncePurchased
- 24/7OnlineChat Support
- ItsLatestVersion

Visit us at: <https://www.certswarrior.com/exam/nurse-practitioner>

Latest Version: 6.0

Question: 1

To promote family integration, the best time for siblings to visit the mother and neonate is

- A. as soon after delivery as possible
- B. just prior to discharge from hospital
- C. after discharge from hospital

Answer: A

Explanation:

Children should visit the mother and infant as soon after delivery as possible. Preparation of siblings prior to the birth can decrease anxiety and sibling rivalry by helping the children feel as though they are participants and are valued. Children should be prepared for physical changes in the mother, changing family dynamics, and infant care. A parent/teacher may use dolls to demonstrate childcare and allow the children to practice holding and caring for the baby. When possible, children should have contact with an infant, such as that of a friend or family member.

Question: 2

When treating a neonate for hypothermia, the air temperature should be increased by approximately

- A. 1 °C every hour until infant stabilizes
- B. 2 °C every hour until infant stabilizes
- C. 3 °C every hour until infant stabilizes

Answer: A

Explanation:

Steps in treatment of hypothermia:

Increase the air temperature by approximately 1°C every hour until the infant's temperature is normal and stable.

Determine if the cause of hypothermia is from an abnormal physiological process in the infant or from environmental conditions.

Avoid rewarming the infant too rapidly because rapid rewarming may result in apnea or hypotension.

Maintain the ambient temperature at 1-1.5°C higher than the infant's temperature.

Warm IV fluids with a blood-warming device prior to infusion.

Closely monitor the infant's blood glucose levels, vital signs, and urinary output.

Question: 3

Which of the following drug(s) should always be included with infant resuscitation equipment?

- A. Prostaglandin and epinephrine
- B. Epinephrine
- C. Surfactant and naloxone

Answer: B

Explanation:

The medication that should always be included as part of the minimum neonatal resuscitation equipment is epinephrine. Epinephrine is the AHA recommended pharmacologic intervention in the case of ineffective CPR in neonatal resuscitation. Naloxone is indicated in the case of respiratory distress in infants with opioid withdraw after the administrations of opioids as pain control in the labor process, but is not required to be included. Surfactant (indicated in respiratory distress, particularly in preterm neonates) and prostaglandins (indicated for cyanotic congenital heart defects) are useful in specific cases, but are also not part of the minimum resuscitation equipment

Question: 4

The relationship between the total loading dose of an administered drug and the serum concentration refers to the

- A. absorption
- B. distribution
- C. clearance

Answer: B

Explanation:

Distribution: The volume of distribution is the relationship between the total loading dose of drug administered and the serum concentration (volume of body fluid required to dissolve the amount of drug found in the serum). Absorption: This relates to the rate at which a drug enters the bloodstream and the amount of drug. Clearance: Elimination pathways (liver, kidney) can become saturated if dose of medications is too high or administration is too frequent. Ideally, a drug concentration should be maintained at a steady state (average).

Question: 5

The electrolytes that are of primary concern in the neonate are

- A. calcium, sodium, and potassium
- B. calcium, chloride, and magnesium
- C. sodium, phosphorous, and potassium

Answer: A

Explanation:

Electrolyte	Normal value	Discussion
Calcium	Cord: 8.2-11.2 mg/dL 0-10 days: 17.6-10.4 mg/dL 11 days-2 years: 9.0-11.0 mg/dL	Hypocalcemia: <7 mg/dL is common with infants that are critically ill, IDM, suffered from asphyxia, or are preterm with very low birth weight.
Sodium	Neonate: 133-146 mEq/L	Hypernatremia: >150 mEq/L usually relates to dehydration, use of Na containing solutions, congenital or acquired reduction in ADH, cerebral palsy, and intracranial hemorrhage. Hyponatremia: <130 mEq/L usually relates to overhydration, renal excretion from diuresis, or SIADH.
Potassium	Neonate: 2.7-5.9 mEq/L	Hyperkalemia: >7 mEq/L may relate to renal failure, acidosis, or adrenal insufficiency. Hypokalemia: <3.5 mEq/L usually relates to excessive GI or renal fluid losses.

Question: 6

The major route in a full-term neonate for rapidly increasing body temperature in response to cold stress is

- A. shivering
- B. non-shivering thermogenesis
- C. vasoconstriction

Answer: B

Explanation:

Non-shivering thermogenesis (NST) is the major route of rapid increase of body temperature in response to cold stress in the term neonate. NST is the oxidation of brown fat to create heat. Brown adipose tissue contains a high concentration of stored triglycerides, a rich capillary network and is controlled by the sympathetic nervous system. When brown fat is metabolized, heat is created. When a cold body temperature is detected, the posterior hypothalamus responds by triggering the adrenal glands to release norepinephrine and the pituitary gland to release thyroxine. Both stimulate NST.

Question: 7

On the APGAR scale, a good score is

- A. ≥ 25
- B. ≥ 27

C. ≥ 210

Answer: B

Explanation:

A total score of 27 is a sign of good health with a perfect score being 10. APGAR is a quick evaluation of a newborn's physical condition to determine if emergency medical care is needed and is administered 1 minute and 5 minutes after birth.

Sign	0	1	2
Appearance (Skin Color)	Cyanosis or pallor over entire body	Normal, except for the extremities	Entire body normal
Pulse (Heart Rate)	Absent	<100 bpm	>100 bpm
Grimace (Reflex Irritability)	Unresponsive	Grimace	Infant sneezes, coughs, and recoils
Activity (Muscle Tone)	Absent	Flexed limbs	Infant moves freely
Respiration (Breathing Rate and Effort)	Absent	Bradypnea, dyspnea	Good breathing and crying

Question: 8

The initial treatment for patent ductus arteriosus (PDA) for a preterm neonate is usually

- A. prostaglandin
- B. surgical ligation
- C. indomethacin or ibuprofen lysine

Answer: C

Explanation:

Patent ductus arteriosus (PDA) is failure of the ductus arteriosus that connects the pulmonary artery and aorta to close after birth, resulting in left to right shunting of blood from the aorta back to the pulmonary artery. This increases the blood flow to the lung and causes an increase in pulmonary hypertension that can result in damage to the lung tissue. Treatment for PDA:

- Indomethacin (Indocin®) or ibuprofen lysine given within 10 days of birth is successful in closing about 80% of defects.
- Surgical repair with ligation of the patent vessel if conservative treatment is unsuccessful.

Question: 9

Insensible water loss in neonates includes

- A. urination
- B. evaporated fluids
- C. stool

Answer: B

Explanation:

Insensible water losses (IWL) occur as water evaporates from the skin (2/3) or the respiratory tract (1/3). IWL cannot be directly measured. Premature neonates have thin skin that allows for increased amounts of evaporative water loss. As the skin matures and the stratum corneum develops (around 31 weeks of gestation) less water is lost through the skin. A full-term neonate will have an IWL of 12/ml/kg/24 hours at 50% humidity. Factors that increase IWL include prematurity, radiant warmers, phototherapy, fever, low humidity, and tachypnea. Sensible water losses occur via urination, stool, and gastric drainage and can be accurately measured.

Question: 10

A neonate, 2 days after birth, develops a generalized rash with erythematous papules, vesicles, and some pustules everywhere but on the palms and soles of feet. The most likely diagnosis is

- A. erythema toxicum
- B. neonatal pustular melanosis
- C. cutis marmorata

Answer: A

Explanation:

Erythema toxicum is a skin eruption of erythematous papules, vesicles, and sometimes pustules. Erythema toxicum is essentially benign and occurs in 250% of newborns. It is a generalized rash everywhere except the palms and soles of the feet, usually occurring 2-3 days after birth. Neonatal pustular melanosis is a benign rash (vesicles and macules) but not associated with erythema. Cutis marmorata is a disorder in which the infants skin mottles or marbles when exposed to cold, because the superficial blood vessels dilate and contract at the same time.



CERTSWARRIOR

FULL PRODUCT INCLUDES:

Money Back Guarantee



Instant Download after Purchase



90 Days Free Updates



PDF Format Digital Download



24/7 Live Chat Support



Latest Syllabus Updates



For More Information – Visit link below:

<https://www.certswarrior.com>

16 USD Discount Coupon Code: U89DY2AQ