



CERTSWARRIOR

# Healthcare CCI-RCIS

**CCI Registered Cardiovascular Invasive Specialist (RCIS)**

**Questions&AnswersPDF**

**ForMoreInformation:**

**<https://www.certswarrior.com/>**

## **Features:**

- 90DaysFreeUpdates
- 30DaysMoneyBackGuarantee
- InstantDownloadOncePurchased
- 24/7OnlineChat Support
- ItsLatestVersion

Exam Code: CCI-RCIS  
Exam Name: CCI Registered Cardiovascular Invasive Specialist (RCIS)  
Exam-Total-Time: 180  
Exam-Version: 6.0  
Exam-Total-Marks: 500  
Vendor-Name: Healthcare  
Total Questions: 110  
RatingbyUsers: 3

## Question: 1

The routine Demerol dosage that is used for conscious sedation would be considered \_\_\_\_\_ mg with a maximum dosage of \_\_\_\_ mg?

- A. 1-2mg/10mg
- B. 10-20mg/100mg
- C. 40-50mg/50mg
- D. None of the above

**Answer: B**

### Explanation:

The routine Demerol dosage used for conscious sedation generally falls within the range of 10-20 milligrams (mg). This dosage is considered effective for inducing a state of sedation where the patient remains awake but is significantly more relaxed and less sensitive to pain. Conscious sedation is commonly used in minor procedures that do not require full anesthesia but still need the patient to be relaxed and somewhat unaware of the procedure, thus easing both patient anxiety and procedural pain. The maximum dosage of Demerol for conscious sedation should not exceed 100 mg. This cap is placed to avoid potential overdose and severe side effects, which can include respiratory depression, severe hypotension, and the risk of increased patient morbidity. It is also important to consider the patient's weight when calculating the maximum dose. A common guideline used by healthcare providers is limiting the dosage to no more than 1.5 mg per kilogram (kg) of the patient's body weight. This means that for a patient weighing 70 kilograms (about 154 pounds), the maximum dose would not exceed 105 mg, aligning well with the 100 mg maximum guideline.

It is crucial for healthcare providers to assess each patient individually, considering factors such as age, body weight, overall health condition, and concurrent medication use. This ensures that the administered dosage is both safe and effective. Monitoring during and after administration of Demerol is also essential to swiftly address any adverse reactions or complications that may arise.

In summary, the routine Demerol dosage for conscious sedation is typically 10-20 mg, with a strict maximum set at 100 mg. This dosage helps manage pain and anxiety during procedures that require the

patient to be sedated but not fully unconscious, ensuring the safety and comfort of the patient throughout the procedure.

## Question: 2

Which of the following would not be considered a part of the pre-procedural assessment for the patient?

- A. A review of all their major organ systems
- B. Review of past bone fractures
- C. History of any drug use
- D. History of smoking

**Answer: B**

Explanation:

The question asks which of the listed items would not typically be included in a pre-procedural assessment for a patient. Out of the options provided, "Review of past bone fractures" stands out as the least relevant to most pre-procedural assessments, unless the procedure specifically relates to orthopedic issues or past fractures may directly impact the procedure or anesthesia.

Pre-procedural assessments are primarily aimed at identifying any potential risks or complications that might arise during or after a medical or surgical procedure. The focus is generally on evaluating the patient's current health status and medical history that are directly relevant to the procedure they are undergoing.

Typical components of a pre-procedural assessment include: - **A review of all their major organ systems**: This is crucial to determine if organs like the heart, lungs, liver, and kidneys are functioning well enough to handle the stress of the procedure and the anesthesia. - **History of any drug use**: Important for identifying potential interactions with anesthesia or other medications used during the procedure. - **History of any alcohol use**: Chronic alcohol use can affect liver function and the patient's response to medications. - **History of any smoking**: Smoking can affect respiratory function and healing times, which are critical considerations for anesthesia and recovery. - **History of any previous problems or experiences with sedative agents**: This helps in planning anesthesia to avoid repeating past complications.

In contrast, a review of past bone fractures is generally not part of this assessment unless the procedure directly involves areas previously affected by fractures or if there is a concern about bone density that might affect surgical outcomes (such as in spinal surgeries). Therefore, for most general procedures, reviewing past bone fractures does not hold as much immediate relevance as the other factors listed.

## Question: 3

When converting pounds to kilograms, which of the following would be correct for converting 298 pounds to kilograms?

- A.  $298 + 2.2 = 300.2$  kg
- B.  $298 \text{ divided by } 2.2 = 135.45$  kg
- C.  $298 \text{ divided by } 4.4 = 67.72$  kg

D. None of the above

**Answer: B**

Explanation:

To convert pounds to kilograms, a specific conversion factor is used. This factor is based on the fact that one kilogram is approximately equal to 2.20462 pounds. Therefore, to convert a weight from pounds to kilograms, the weight in pounds is divided by 2.20462.

In your question, the options given explore different computations involving the number 2.2, which is a rounded form of the more precise 2.20462. The simplification to 2.2 makes the math easier while still providing a reasonably accurate conversion.

The correct method to convert pounds to kilograms, as per your question, is by dividing the weight in pounds by 2.2. This is because 2.2 is a practical, although slightly less precise, approximation of 2.20462, making it a commonly used value in everyday conversions that do not require high precision.

For the specific example given in the question, converting 298 pounds to kilograms using the approximate conversion factor of 2.2 would be calculated as follows:

$$\text{Kilograms} = \text{Pounds} / 2.2 = 298 / 2.2 = 135.45 \text{ kg}$$

This calculation shows that 298 pounds is approximately equal to 135.45 kilograms when using the simplified conversion factor. Therefore, the correct choice among the provided options would be:

298" divided by " 2.2=135.45" kg"

Additionally, the explanation provided in the question, stating that the formula for converting pounds to kilograms is "Pounds divided by 2.2 = kilograms," is accurate and aligns with widely used conversion practices.

### Question: 4

An advantage for using the power injector when administering contrast medium during a catheterization procedure would be considered all of the following except?

- A. Ensures the automatic recording of the injection.
- B. Ensures that the amount of the contrast medium being injected would be the correct volume.
- C. Ensures the patient will have no complications.
- D. Ensures that the medium is injected at a constant pressure.

**Answer: C**

Explanation:

The question asks which option is NOT an advantage of using a power injector when administering contrast medium during a catheterization procedure. To answer this, we need to examine each provided choice against the known benefits of using a power injector.

Firstly, power injectors are designed to enhance the delivery of contrast media during imaging procedures. One significant advantage is the ability to ensure that the contrast medium is injected at a consistent rate and pressure. This is crucial for obtaining high-quality images, as inconsistent pressure can lead to poor image quality and the potential need for repeat scans, which increases the patient's exposure to radiation and contrast media.

Another advantage of using power injectors is the ability to deliver precise volumes of contrast media. This precision ensures that the exact, pre-determined amount of contrast needed to achieve optimal imaging results is administered, reducing the risk of under- or over-dosing which can affect image quality and patient safety.

Furthermore, power injectors facilitate the automatic recording of all injection parameters. This includes the amount of contrast injected, the rate of injection, and the total volume used. This automated documentation is vital for maintaining accurate medical records and can be helpful for reviewing the procedure's details if there are any concerns about patient outcomes or if further analysis is required. Now, evaluating the options: 1. Ensures the automatic recording of the injection - True, as discussed, this is a benefit. 2. Ensures the patient will have no complications - This statement is misleading. While using a power injector can help reduce certain risks associated with manual injections, such as human error in dosing, it does not guarantee that the patient will have no complications. Complications can arise due to factors unrelated to the method of injection, such as allergic reactions to the contrast medium or other patient-specific issues. 3. Ensures that the amount of the contrast medium being injected would be the correct volume - True, as power injectors are specifically designed to provide this accuracy. 4. Ensures that the medium is injected at a constant pressure - True, a fundamental benefit of using power injectors.

Therefore, the correct answer to the question, "An advantage for using the power injector when administering contrast medium during a catheterization procedure would be considered all of the following except?" is "Ensures the patient will have no complications." This choice is the only one that does not directly relate to the specific capabilities and advantages of using a power injector.

### Question: 5

Which of the following measurements are typically taken in the inferior vena cava during a right heart catheterization?

- A. Pulmonary capillary wedge pressure
- B. O2 saturation
- C. PA pressure
- D. All of the above

**Answer: B**

Explanation:

During a right heart catheterization, a procedure primarily used to diagnose and evaluate conditions related to the heart and lungs, various hemodynamic parameters are measured. This procedure involves the insertion of a catheter into the right side of the heart and often into the pulmonary artery. One of the key measurements taken during this procedure is the oxygen saturation (O2 saturation) level in different parts of the circulatory system.

The inferior vena cava, which is a large vein that carries deoxygenated blood from the lower and middle parts of the body to the heart, is one of the sites where O2 saturation is measured. The importance of measuring O2 saturation here lies in assessing how well oxygen-depleted blood is being returned to the heart, which is critical for evaluating overall cardiac function and detecting potential abnormalities in the body's oxygen utilization.

In addition to the inferior vena cava, O2 saturation is also measured in other areas during a right heart catheterization. These areas include the superior vena cava, which brings deoxygenated blood from the

upper part of the body; the right atrium, where the blood enters the heart; and the pulmonary artery, which transports blood from the heart to the lungs for oxygenation. By comparing O2 saturation levels across these different sites, physicians can gain detailed insights into the cardiac output and pulmonary function.

However, it's important to note that not all measurements listed, such as pulmonary capillary wedge pressure (PCWP) and pulmonary artery (PA) pressure, are typically measured in the inferior vena cava itself. PCWP, for instance, is measured in the pulmonary artery to estimate left atrial pressure indirectly, and PA pressure is directly measured in the pulmonary artery to assess pulmonary hypertension and other conditions.

Therefore, among the options provided, the measurement typically taken in the inferior vena cava during a right heart catheterization is the O2 saturation. Other parameters like PCWP and PA pressure are measured at different sites in the circulatory system relevant to their specific diagnostic purposes.

### Question: 6

When performing a left ventriculogram, which of the following is typically performed prior?

- A. Aortography
- B. Right ventriculography
- C. Right heart catheterization
- D. Coronary angiogram

**Answer: D**

Explanation:

In the context of cardiac diagnostic procedures, a left ventriculogram is a specialized X-ray test used to assess the structure and function of the left ventricle, which is the heart's main pumping chamber. This test provides crucial information about the volume of the chamber, the condition of the heart valves, and the movement of the heart wall.

Before performing a left ventriculogram, it is common practice to conduct a coronary angiogram. A coronary angiogram is a procedure that uses X-ray imaging to see your heart's blood vessels. The purpose of performing a coronary angiogram before a left ventriculogram is multifold: 1. **\*\*Assessment of Coronary Arteries:\*\*** The coronary angiogram provides a detailed view of the coronary arteries and the blood flow within them. This is vital for detecting any blockages or narrowing that could affect the heart's function, contributing to conditions like ischemia or heart attacks which directly influence the results and interpretation of the left ventriculogram. 2. **\*\*Safety Considerations:\*\*** Knowing the status of the coronary arteries helps the medical team to assess the risk of inducing myocardial ischemia or other complications during the left ventriculogram. If severe coronary artery disease is present, the approach to the left ventriculogram might be adjusted to minimize stress on the heart. 3.

**\*\*Comprehensive Evaluation:\*\*** Performing the coronary angiogram first allows physicians to gather comprehensive data about the patient's heart health. This information can guide subsequent interventions and treatments, including decisions made during the left ventriculogram regarding potential valve repairs or assessments of heart muscle viability.

In conclusion, while both aortography, right ventriculography, and right heart catheterization are valuable procedures in specific contexts, the coronary angiogram is particularly critical before a left ventriculogram. It serves to ensure the safety of the patient while providing essential diagnostic data that can influence the course of treatment, including surgical decisions or the need for further

interventions like stenting or bypass surgery. Thus, the coronary angiogram is not only a preparatory step but a foundational procedure that enhances the diagnostic accuracy and safety of the left ventriculogram.

### Question: 7

What color would the chronic effusion fluid typically be?

- A. Red
- B. Yellow
- C. Black
- D. Brown

**Answer: B**

Explanation:

The color of chronic effusion fluid is typically a clear yellow. This coloration is primarily due to the presence of protein and other cellular elements that have been filtered from the blood into the effusion fluid over a prolonged period. Chronic effusions generally indicate a slow, ongoing process where the body is reacting to inflammation, irritation, or low-grade injury.

However, the appearance of the effusion fluid can vary depending on the underlying cause and the duration of the effusion. In some cases, the effusion may appear serosanguineous, which is a mixture of clear yellow fluid and blood, giving the fluid a pinkish tinge. This can occur if there is intermittent bleeding into the effusion space, which could be due to mild ongoing trauma or irritation in the affected area.

In other instances, chronic effusion fluid may turn dark brown. This change in color typically occurs when there is repeated bleeding over a long period, and the blood cells in the effusion degrade. The hemoglobin from the red blood cells breaks down into hemosiderin and other pigments, which can give the fluid a dark brown color, often described as resembling tea or cola.

It is important to contrast these with acute effusions, which are usually due to more sudden causes like trauma, infections, or artery perforation. The fluid in acute effusions is often frankly bloody or very red, reflecting the active and recent nature of blood leakage into the space.

In summary, while the typical color of chronic effusion is clear yellow, variations like serosanguineous or dark brown can also occur depending on specific conditions and the dynamics of the effusion. Each color variation provides clues about the nature of the underlying pathological process and can guide further diagnostic and therapeutic approaches.

### Question: 8

The most common congenital cardiac abnormality would be considered which of the following?

- A. Ventricular septal defect
- B. Atrial septal defect
- C. Tetralogy of Fallot
- D. AV defect

**Answer: A**

Explanation:

The most common congenital cardiac abnormality is the ventricular septal defect (VSD). A VSD is characterized by an opening or a hole in the septum that separates the heart's two lower chambers, known as the ventricles. This defect allows blood to pass from the left to the right ventricle, bypassing the normal route of circulation, which can lead to various complications.

The occurrence of VSD can vary in size and impact. Small VSDs may not cause significant problems and often close on their own as the child grows. However, larger VSDs can lead to serious cardiac issues. These include increased pulmonary blood flow, which can strain the heart and lungs, potentially leading to congestive heart failure, pulmonary hypertension, or Eisenmenger's syndrome if left untreated. Treatment for VSD depends on the size and severity of the defect, as well as the presence of symptoms. Monitoring is often recommended for smaller defects to observe if spontaneous closure occurs. For larger or symptomatic VSDs, treatment options include surgical repair or catheter-based procedures. Surgical correction involves placing a patch or stitches to close the defect, thereby preventing the abnormal flow of blood between the ventricles. This is usually performed via open-heart surgery, which has a high success rate and generally good outcomes for most children.

In conclusion, among congenital heart defects, the ventricular septal defect stands out as the most common. Understanding the nature of this defect, its potential complications, and treatment options is crucial for effective management and improving the quality of life for affected individuals. Early detection and appropriate intervention can significantly influence the prognosis and allow for a relatively normal life for many children with a VSD.

## Question: 9

Which of the following might be considered ways to help avoid the development of air embolisms during a cardiac catheterization?

- A. Check any syringes for potential air pockets
- B. Carefully back bleed the lines.
- C. Careful flushing of all connections.
- D. All of the above

**Answer: D**

Explanation:

In the context of cardiac catheterization, an air embolism—a dangerous condition where air bubbles enter the bloodstream and obstruct blood flow—can pose significant risks to the patient. To mitigate this risk, there are several precautionary techniques that can be employed by medical professionals to ensure the procedure is as safe as possible. Below is an expansion on each of the mentioned strategies:

**\*\*Checking any syringes for potential air pockets:\*\*** Before any syringe is used during a cardiac catheterization, it is crucial to check for and remove any air pockets. This is because the introduction of air into the vascular system through syringes is a common way that air embolisms can occur. Healthcare providers are trained to expel any air from a syringe by pointing it upright and tapping its side, allowing the air bubbles to rise to the top and be expelled before the syringe's contents are administered.



**\*\*Carefully back bleeding the lines:\*\*** This technique involves allowing a small amount of blood to flow back into the catheter or line after it is inserted, but before it is locked in place and used for the introduction of any medications or contrast agents. This back bleeding helps to ensure that any air that might have been introduced during the insertion of the catheter is expelled.

**\*\*Careful flushing of all connections:\*\*** This preventive measure involves the thorough flushing of all catheters and connections with saline or another appropriate solution before they are used. This process helps to push out any air that may be present in the lines or connections used during the procedure.

By consistently implementing these practices, the risk of air embolisms during cardiac catheterizations can be significantly reduced, thereby enhancing the safety of the procedure for patients. It is essential for medical personnel to be meticulous in their approach and adhere strictly to these preventive measures.

### Question: 10

Which of the following medications have the potential to impact the risks for a cardiac catheterization?

- A. Ibuprofen
- B. Warfarin
- C. Tylenol
- D. Both A and B

**Answer: D**

Explanation:

Cardiac catheterization is a medical procedure used to diagnose and treat certain cardiovascular conditions. During this procedure, a long thin tube called a catheter is inserted in an artery or vein in your groin, neck, or arm and threaded through your blood vessels to your heart. Given the invasive nature of the procedure, certain medications can affect the risks associated with cardiac catheterization. To address the question of which medications might impact the risks of cardiac catheterization, we need to consider how these drugs interact with the body's cardiovascular and coagulation systems, among other factors. The choices given include Ibuprofen, Warfarin, Tylenol (Acetaminophen), and an option indicating both Ibuprofen and Warfarin (Both A and B).

1. **\*\*Ibuprofen\*\***: This is a nonsteroidal anti-inflammatory drug (NSAID). NSAIDs are known to potentially affect blood clotting, which can increase the risk of bleeding during and after the procedure. They may also affect kidney function, which is crucial in the context of cardiac catheterization where contrast agents are used that can impact kidney function.

2. **\*\*Warfarin\*\***: Warfarin is an anticoagulant, or a blood thinner, which is used to prevent blood clots. While beneficial in preventing strokes or other clot-related issues, its use can substantially increase the risk of bleeding during invasive procedures like cardiac catheterization. Management of patients on Warfarin undergoing such procedures typically involves careful monitoring of blood coagulation levels and possibly adjusting the dosage or temporarily stopping the medication.

3. **\*\*Tylenol (Acetaminophen)\*\***: Unlike NSAIDs and Warfarin, Tylenol does not typically impact blood clotting in significant ways under normal circumstances, nor does it have the same level of renal interactions. Therefore, it is less likely to affect the risks associated with cardiac catheterization concerning bleeding or kidney function.

4. **\*\*Both A and B (Ibuprofen and Warfarin)\*\***: Choosing this option recognizes that both Ibuprofen and Warfarin have potential impacts on the risks associated with cardiac catheterization, primarily due to their effects on blood clotting and, in the case of Ibuprofen, possibly kidney function.

In conclusion, while Tylenol is generally safer in the context of not affecting bleeding or kidney function significantly, both Ibuprofen and Warfarin carry increased risks during cardiac catheterization. The option "Both A and B" correctly identifies that both these medications (Ibuprofen and Warfarin) can alter the risk profile for this procedure, mainly due to their impact on coagulation and the potential for bleeding complications.



# CERTSWARRIOR

## *FULL PRODUCT INCLUDES:*

Money Back Guarantee



Instant Download after Purchase



90 Days Free Updates



PDF Format Digital Download



24/7 Live Chat Support



Latest Syllabus Updates



For More Information – Visit link below:

**<https://www.certswarrior.com>**

**16 USD Discount Coupon Code: U89DY2AQ**