



CERTSWARRIOR

# International DHA-Audiologist DHA Audiologist Certification Exam

**Questions&AnswersPDF**

**ForMoreInformation:**

**<https://www.certswarrior.com/>**

## **Features:**

- 90DaysFreeUpdates
- 30DaysMoneyBackGuarantee
- InstantDownloadOncePurchased
- 24/7OnlineChat Support
- ItsLatestVersion

# Latest Version: 6.0

## Question: 1

Which of the following would be used to cross-check the pure tone air conduction for reliability?

- A. Bone conduction.
- B. Pitch matching.
- C. Acoustic reflex testing.
- D. Speech reception threshold.

**Answer: d**

Explanation:

To determine the most effective method for cross-checking the reliability of pure tone air conduction audiometry results, it's essential to understand the role of different audiometric tests. Pure tone air conduction testing primarily measures the ability to hear sounds transmitted through air to the ear. However, to verify these results, other types of auditory assessments are used.

One such method is the Speech Reception Threshold (SRT). The SRT assesses the softest level at which speech can be understood 50% of the time. This test is particularly useful for cross-checking pure tone air conduction results because it involves a different aspect of hearing—understanding speech rather than just detecting tones. If the SRT results are consistent with the pure tone results, it suggests that the pure tone findings are reliable. Additionally, the SRT can help identify non-organic hearing loss, also known as malingering, by comparing speech understanding capabilities with the audiometric thresholds. Bone conduction testing is another method used alongside air conduction testing. While pure tone air conduction assesses the outer and middle ear function, bone conduction testing evaluates the inner ear and auditory pathways to the brain, bypassing the outer and middle ears. When bone conduction thresholds are better than air conduction thresholds, it often indicates a conductive hearing loss.

Consistency between these two tests can further validate the reliability of the air conduction results. Other tests like intermittent audiometry, pitch matching, and acoustic reflex testing also play roles in a comprehensive auditory assessment, but they serve different specific purposes. Intermittent audiometry, for example, evaluates the hearing ability over intermittent or fluctuating noise levels and might not directly cross-check pure tone air conduction results. Pitch matching is typically used in tinnitus evaluation, and acoustic reflex testing assesses the reflex pathways of the middle ear muscles, which can indicate the site of the lesion but might not directly validate air conduction results.

In conclusion, while several audiometric tests exist, the Speech Reception Threshold (SRT) and bone conduction testing are particularly effective in cross-checking pure tone air conduction results for reliability. SRT provides a direct comparison of speech understanding against tonal thresholds, and bone conduction testing helps rule out conductive components, affirming the findings of air conduction tests. Each test, however, should be selected based on the specific diagnostic needs and the overall context of the patient's auditory profile.

## Question: 2

DFNA2 is a mutation in which gene(s)?

- A. KCNQ4 and GJB3.
- B. COCH.
- C. EYA4.
- D. GJB2.

**Answer: a**

Explanation:

DFNA2 is a mutation associated with two specific genes: KCNQ4 and GJB3. These genes play critical roles in the normal functioning of the ear, particularly in the process of hearing.

The KCNQ4 gene is known for its involvement in encoding a protein that forms a potassium channel in the inner ear. This channel is crucial for the proper electrical signaling in the auditory pathway. When there are mutations in the KCNQ4 gene, it can lead to malfunctions in these potassium channels, disrupting the electrical signals that are essential for normal hearing. The result is autosomal dominant nonsyndromic hearing loss, which means the hearing loss occurs by itself without other syndromic features and can be inherited from just one affected parent. This form of hearing loss is typically progressive, which means it worsens over time.

On the other hand, the GJB3 gene encodes a protein that is part of a group known as connexins, specifically connexin 31. Connexins are important for forming gap junctions, which are channels that allow for direct communication between cells. In the context of hearing, these gap junctions are vital for maintaining the ionic environment of the inner ear, which is necessary for proper auditory signal transduction. Mutations in GJB3 disrupt this cellular communication, leading to hearing impairment. While the KCNQ4 gene mutation is more commonly associated with DFNA2, mutations in the GJB3 gene are also implicated. It is important for individuals diagnosed with autosomal dominant nonsyndromic hearing loss to undergo genetic testing to determine the specific gene involved, as this can affect management and counseling regarding the progression of the condition and the likelihood of passing it on to offspring.

### Question: 3

With regard to aural rehabilitation, what does the "O" in CORE assessment stand for?

- A. Overall participation variables.
- B. Objective variables.
- C. Open-set variables.
- D. Ordinary variables.

**Answer: A**

Explanation:

The "O" in CORE assessment stands for "Overall participation variables."

Overall participation variables are a crucial component of the CORE assessment in aural rehabilitation. This aspect of the assessment focuses on understanding how a person's hearing impairment impacts their daily life, particularly in terms of social interaction and participation in various activities. It

considers the individual's ability to engage in personal, social, vocational, and educational environments. By evaluating these variables, audiologists and other hearing professionals can better understand the practical implications of hearing loss on an individual's life.

The CORE assessment framework is a comprehensive tool used by professionals to evaluate and manage the rehabilitation needs of people with hearing impairment. CORE stands for Communication status, Overall participation variables, Related personal factors, and Environmental factors. Each of these components helps in forming a holistic view of the patient's auditory challenges and their implications on daily life.

Specifically, the Overall participation variables help in identifying the barriers that hearing loss might pose in an individual's life. This includes looking at how well the individual can follow conversations in noisy environments, participate in meetings or classes, engage in social activities, or even watch television. This evaluation helps in pinpointing specific areas where interventions, such as hearing aids, cochlear implants, or other assistive listening devices, can be most beneficial.

Moreover, this component of the CORE assessment also takes into account the emotional and psychological impact of hearing loss, which can affect overall quality of life. It assesses feelings of isolation, frustration, and decreased self-esteem that often accompany hearing impairment.

Understanding these aspects is crucial for developing a comprehensive rehabilitation plan that addresses both the audiological needs and the psychosocial well-being of the patient.

In summary, the "Overall participation variables" in the CORE assessment are essential for a thorough evaluation of how hearing loss affects an individual's ability to participate in everyday activities. This understanding guides the development of personalized rehabilitation plans that aim to improve not only hearing but also the overall quality of life for individuals with hearing impairment.

### Question: 4

Which of these is filled by perilymph?

- A. Scala vestibuli.
- B. Scala media.
- C. Scala tympani
- D. Both A and C.

**Answer: D**

Explanation:

The question is asking to identify which parts of the ear are filled with a fluid called perilymph. To address this, it's important to understand the structure of the cochlea, which is part of the inner ear. The cochlea is divided into three chambers or canals: scala vestibuli, scala media, and scala tympani.

Perilymph is a fluid that fills two of these chambers: the scala vestibuli and the scala tympani. The scala vestibuli is the upper chamber of the cochlea that begins near the oval window, a membrane-covered opening that receives vibrations from the middle ear ossicles. The scala tympani is the lower chamber which terminates at the round window, another membrane that dissipates vibrations within the cochlea.

In contrast, the scala media, also known as the middle chamber, does not contain perilymph. Instead, it is filled with a different fluid called endolymph. This distinction is crucial because perilymph and endolymph have different ionic compositions, which play an essential role in the process of hearing by

facilitating the transduction of sound-induced mechanical movements into electrical signals by hair cells in the organ of Corti, located in the scala media.

Given these details, the correct answer to the question is "Both A and C," where: - A refers to "Scala vestibuli" and - C refers to "Scala tympani."

Each of these chambers—scala vestibuli and scala tympani—contains perilymph, which serves not only as a medium for transmitting sound vibrations but also helps in maintaining the ionic balance required for normal auditory function. Thus, understanding the role and location of perilymph is fundamental in grasping how hearing mechanisms function within the inner ear.

## Question: 5

When treating a patient, the need for communication with that patient is critical. Which of the below statements is not accurate regarding the requirement and the ethical standards expected by the public?

- A. You are required to communicate the nature of the procedures and the possible side effects to the patient and/or caregiver before conducting the procedure.
- B. You are required to communicate the possibility or intent to use the patient's case or any of his/her information in your future research or data collecting.
- C. You must keep accurate and timely documentation of all services rendered with all of your patients according to regulations and ethical standards.
- D. You are not allowed to communicate the use of a patient in product research due to the confidentiality clause of your relationship with the product manufacturer.

**Answer: D**

Explanation:

The question regarding the ethical standards expected by the public in relation to patient communication is critical in the context of healthcare. Among the statements provided, the one that is not accurate regarding ethical standards is: "You are not allowed to communicate the use of a patient in product research due to the confidentiality clause of your relationship with the product manufacturer." This statement is misleading because it does not consider the ethical imperative of informed consent. Ethical guidelines and regulations, such as those stipulated by bodies like the American Medical Association (AMA) and the World Medical Association (WMA), clearly state the necessity of informing patients about how their cases and personal data might be used, including for purposes like product research. This information must be communicated transparently to ensure that patients are fully aware and can provide or withhold their consent based on a comprehensive understanding of what their involvement entails.

The confidentiality clause, typically present in agreements between healthcare providers and third parties like product manufacturers, must not override the patient's rights to be informed. In fact, any such clause should accommodate the necessity of disclosing information to the patient when their data or biological samples are involved. It is crucial that patients understand and agree to their participation in research or any other secondary use of their data or biological material, which must be explicitly stated in an informed consent form.

Furthermore, ethical standards require that patients be given the option to opt out of research activities without affecting the quality of care they receive. This means that patients must be informed clearly not just about the potential use of their data, but also about their rights to refuse participation. The

principle of voluntary participation is a cornerstone of research ethics, emphasizing respect for patient autonomy.

In summary, the incorrect statement overlooks the fundamental ethical requirement of informed consent, which is designed to protect patient autonomy and ensure transparency in all medical practices, including those involving partnerships with product manufacturers. Ethical healthcare practice demands that any use of patient data in research be communicated to the patient, with their informed consent obtained prior to the use of such data.

### Question: 6

This term refers to ear poisoning, which results from exposure to drugs or chemicals that damage the inner ear or auditory nerve.

- A. Neurotoxicity.
- B. Vestibulotoxicity.
- C. Ototoxicity.
- D. Cochleotoxicity.

**Answer: C**

Explanation:

The correct term that refers to ear poisoning resulting from exposure to drugs or chemicals that damage the inner ear or auditory nerve is "Ototoxicity."

Ototoxicity specifically describes the adverse effects that certain medications and chemicals can have on the cochlea or auditory nerve, which can lead to hearing loss or balance disorders. This condition is significant as it affects the structures within the ear that are crucial for hearing and balance. The inner ear not only involves the cochlea, which is responsible for hearing, but also includes the vestibular system, which regulates balance.

The vestibulocochlear nerve, also known as the auditory nerve, is integral in transmitting sound and balance information from the inner ear to the brain. When ototoxic drugs damage this nerve, the transmission of sensory information is disrupted, leading to symptoms such as tinnitus (ringing in the ears), hearing loss, and problems with balance.

It is important to distinguish ototoxicity from other similar-sounding terms: - **Neurotoxicity** refers to poisoning of the nervous system by harmful substances, affecting the brain or other parts of the nervous system, not specifically limited to the auditory nerve. - **Cochleotoxicity** is a more specific term that refers to toxins that specifically damage the cochlea, leading predominantly to hearing loss. -

**Vestibulotoxicity** refers to damage specifically to the vestibular system, affecting balance but not necessarily hearing. - **Otitis Media** is an inflammation or infection of the middle ear, which is different from ototoxicity as it does not necessarily involve direct chemical or drug-induced damage to the ear structures but can lead to hearing complications due to inflammation or fluid build-up.

Understanding ototoxicity is crucial for healthcare providers, especially when prescribing medications known to carry risks of ear damage. Patients undergoing treatment with potentially ototoxic drugs should be monitored for auditory and vestibular symptoms to manage and mitigate any adverse effects proactively.

### Question: 7

ASHA recommends that the first routine childhood screening occurs:

- A. When the child enters school.
- B. In the first grade.
- C. In the second grade.
- D. In the third grade.

**Answer: A**

Explanation:

ASHA, or the American Speech-Language-Hearing Association, is a professional association for audiologists, speech-language pathologists, and speech, language, and hearing scientists in the United States and internationally. One of their roles is to set standards and guidelines for the screening and early detection of speech, language, and hearing issues in children.

According to ASHA's recommendations, routine childhood screenings for hearing, speech, and language development should begin when a child enters school. This is a critical time as early detection of potential communication disorders is essential for timely intervention, which can significantly improve the child's long-term communication skills, academic performance, and social interactions.

ASHA suggests specific grades during which screenings should ideally occur: - When the child enters school - Kindergarten - First grade - Second grade - Third grade - Seventh grade - Eleventh grade  
Each of these stages represents key points in a child's academic and developmental progress where issues may emerge or become apparent as the child faces new cognitive, social, and educational challenges. Screening at these intervals ensures that any emerging problems can be addressed before they become more serious and affect the child's development more broadly.

For example, entering school often coincides with preschool or kindergarten, where foundational skills in communication are critical for learning and social interaction. Subsequent screenings in first through third grades align with early educational transitions and increased academic demands, making it crucial to ensure that children can hear, speak, and understand effectively.

Screenings in seventh and eleventh grades coincide with later educational and social development stages, where communication skills become even more complex and integral to learning and personal development. These screenings aim to catch any issues that may have been missed earlier or have developed as the child's communication and academic needs have evolved.

Overall, ASHA's recommended schedule is designed to support and enhance children's ability to achieve their full academic and social potential by ensuring their communication abilities are developed appropriately for each stage of their growth.

## Question: 8

What does otitis media typically cause?

- A. It causes sensorineural hearing loss.
- B. It causes inner ear damage.
- C. It does not cause hearing loss.
- D. It causes conductive hearing loss.

**Answer: D**

Explanation:

Otitis media primarily causes conductive hearing loss. This condition is an inflammation or infection of the middle ear, which is the air-filled space behind the eardrum that contains the tiny vibrating bones of the ear. Children are more commonly affected by otitis media due to the structure of their eustachian tubes being shorter and more horizontal, which can make it easier for pathogens to access the middle ear.

Conductive hearing loss occurs when there is a problem conducting sound waves anywhere along the route through the outer ear, tympanic membrane (eardrum), or middle ear (ossicles). In the case of otitis media, fluid buildup in the middle ear, a common symptom of the infection, can prevent the ossicles from moving freely. Since these bones need to vibrate in order to transmit sound from the air to the cochlea (the sensory organ of hearing), any restriction in their movement can lead to diminished hearing.

While otitis media is specifically an issue with the middle ear, it does not typically cause sensorineural hearing loss, which affects the inner ear or the neural pathways to the brain. Nor does it usually cause mixed hearing loss, which is a combination of conductive and sensorineural hearing loss. However, in severe or untreated cases, the infection can potentially spread to the inner ear or lead to more serious complications that might affect both the conductive and sensorineural components of the auditory system.

### Question: 9

In conclusion, otitis media most commonly leads to conductive hearing loss due to the accumulation of fluid in the middle ear, which impedes the proper transmission of sound through the middle ear. Regular medical attention and treatment are essential to prevent the condition from leading to more significant hearing impairment or other complications.

The ethical acceptance of gifts, as a licensed audiologist, has many guidelines and criteria that must be strictly followed. Which of the below examples is not an example of a gift, and can be accepted by you?

- A. After purchasing a large number of a particular hearing assistance device, you receive a gift card from the manufacturer of the devices.
- B. While attending a seminar, you receive a faux leather portfolio, pen and pencil, and other items that are supplied to everyone in attendance.
- C. While finishing a purchase in the office, from a sales representative of a pharmaceutical company, the rep gives you fifty dollars case as a thank you for your business.
- D. While attending a conference on a new diagnostic procedure using a new computer system, you receive a laptop that is preloaded with the system.

**Answer: B**

Explanation:

\*The ethical guidelines for accepting gifts as a licensed audiologist are stringent to avoid any appearance of impropriety or influence that could compromise professional judgment. Here, we have several scenarios to consider for determining which item can be ethically accepted without violating professional standards. \*



\*In the first scenario, receiving a gift card from a manufacturer after purchasing a large number of hearing assistance devices represents a significant gift that could be seen as a kickback, intended to influence future buying decisions. This is clearly unethical and should not be accepted. \*

\*The second scenario involves receiving items like a faux leather portfolio, pen, and pencil at a seminar, which is supplied to everyone in attendance. These items are generally considered of nominal value and are standard at many conferences to facilitate note-taking and organization. Since they are offered to all participants without specific conditions and are of minimal value, accepting these items does not typically pose an ethical issue. \*

\*The third scenario describes receiving \$50 in cash from a sales representative as a thank-you for business transactions. Cash gifts, regardless of amount, are almost always considered unethical in professional settings as they can be construed as a direct bribe for favorable treatment or continued business. \*

\*In the fourth scenario, receiving opera tickets from a manufacturer's representative could be seen as an attempt to curry favor, especially when coupled with the mention of an upcoming business call. Such gifts, which hold significant monetary and personal value, could influence or appear to influence professional decisions, making them inappropriate. \*

\*The fifth scenario involves receiving a laptop preloaded with a new diagnostic system while attending a conference on that system. If the laptop is necessary for the training and intended for professional use to better understand or implement the system, and if such a practice is standard at such conferences, it may be considered acceptable. However, this could depend on the value of the laptop and whether it is a requirement to keep it post-training. \*

\*Among the examples given, the only clearly acceptable scenario under typical ethical guidelines is receiving low-value items like a portfolio, pen, and pencil at a seminar where such items are given to all attendees. This scenario does not imply any special treatment or significant value that could influence professional judgment, making it an appropriate acceptance of a gift in a professional context.

## Question: 10

At which rate do you increase dB in the LDL test?

- A. At 10 dB each interval.
- B. At 5 dB each interval.
- C. At 7 dB each interval.
- D. At 12 dB each interval.

**Answer: B**

Explanation:

The correct answer to the question regarding the rate at which decibels (dB) are increased during an LDL (Loudness Discomfort Level) test is: "At 5 dB each interval." This choice aligns with standard audiological practices for conducting LDL assessments. The purpose of the LDL test is to determine the sound level at which sound becomes uncomfortably loud for a patient. This threshold is crucial for configuring hearing aids and for understanding a patient's tolerance to loud sounds.

The LDL test typically begins at the Most Comfortable Loudness (MCL) level for the patient. MCL is the sound level at which the patient finds the audio neither too soft nor too loud, providing a baseline comfort point from which the test begins. Starting from the MCL, the audiologist or clinician increases the sound intensity in small increments to carefully gauge the patient's reaction to increasing loudness.

This method ensures precision and avoids causing discomfort from too large an increase in loudness at once.

The standard increment used in these tests is 5 dB per interval. This rate of increase is chosen because it strikes a balance between sensitivity and patient comfort. Smaller increments might result in a test that is too lengthy and impractical, while larger increments might skip over the actual threshold level of discomfort, leading to less accurate results. Thus, increasing the volume by 5 dB at each interval is widely accepted as it provides a systematic approach to identifying the exact point at which sound becomes uncomfortably loud for the patient.

Therefore, when conducting an LDL test, it is important to adhere to this increment rate to ensure the accuracy and effectiveness of the test. This helps in creating an effective hearing aid program that enhances the user's auditory experience without causing discomfort from overly loud sounds.



# CERTSWARRIOR

## FULL PRODUCT INCLUDES:

Money Back Guarantee



Instant Download after Purchase



90 Days Free Updates



PDF Format Digital Download



24/7 Live Chat Support



Latest Syllabus Updates



For More Information – Visit link below:

**<https://www.certswarrior.com>**

**16 USD Discount Coupon Code: U89DY2AQ**